

Ad ♂, SA:60 Ad ♀, SA:61 Juv ♂). Both adults were seen feeding the juvenile on separate occasions, clearly indicating recent local breeding, although the testes of the male had already regressed. Their plumage is very worn, with no patterning left on the feathers, although the female has started moulting the inner 2 primaries. Unfortunately the specimens were all somewhat damaged during initial storage in a freezer, but the juvenile plumage can be partially described:- upperparts, wings and tail very dark brown with large buff spots on the back and head; rufous-buff tips to the greater coverts forming a narrow wing bar; secondaries narrowly edged with rufous, the tertials with broad buff edges; the tail with a narrow rufous tip. From what is left of the feathers, the underparts appear similar to those of the adult, but the chest markings are darker brown.

Acknowledgements. CMF wishes to thank Mr. Patrick Andau, Assistant Chief Game Warden of Sabah, for support of ornithological research in the State and for cooperation in allowing him to conduct a bird-ringing programme in Sepilok forest reserve. F. H. Sheldon and J. Kennard who organised collecting on behalf of the Western Foundation of Vertebrate Zoology and L. F. Kiff, Director of the Foundation, very kindly allowed DRW to review their *Muscicapa* material, now all held in Los Angeles.

References:

- Amadon, D. & duPont, J. E. 1970. Notes on Philippine birds. *Nemouria* 1: 1-14.
 Siebers, H. C. 1928. Neue vögel von Sumba. *Treubia* 10: 399-404.
 Wells, D. R. 1977. *Muscicapa williamsoni* Deignan: a reappraisal. *Bull. Brit. Orn. Cl.* 97: 83-87.
 — 1982. Notes on some representatives of the Brown Flycatcher *Muscicapa latirostris* Raffles in Southeast Asia. *Bull. Brit. Orn. Cl.* 102: 148-153.

Address: D. R. Wells, Zoology Department, University of Malaya, Kuala Lumpur, Malaysia.

© British Ornithologists' Club 1984.

Weights and gonad condition of some Thai birds

by David S. Melville and Philip D. Round

Received 14 March 1984

Many studies on the distribution and taxonomy of birds in Thailand have recorded standard body measurements taken from skins (eg. Chasen & Kloss 1932, Deignan 1945, Riley 1938), but little appears to have been published on the weights of Thai birds. Even though some 185,000 birds of 491 species were ringed in Thailand during the Migratory Animal Pathological Survey 1963-71 (McClure 1974a), the only published weights are those given by McClure & Kwanyuen (1973) for 66 species. Even elsewhere in Southeast Asia, only a very few detailed studies on individual species have been reported (Medway 1973, Nisbet 1967, Nisbet & Medway 1972, Ward 1969). Comparatively little has been published on the seasonality of breeding among birds in Thailand (Herbert 1923-26, McClure 1974b, Round 1982), while the only information on gonad condition is that given by Deignan (1945).

In this paper, we present weight and, in some cases, gonad data for 1686 birds of 165 species, collected or examined alive and subsequently released, from 12 sites in NW, NE, Central and SE Thailand during September 1980 to December 1982. No data were collected during the months June to August, but there is a fairly even spread for the rest of the year, with most data for January, April and December. Details for shorebirds will be presented elsewhere. Details of the study sites are given in Table 1.

TABLE I

Study areas in Thailand for trapping and collecting birds



| Reference | Location | Coordinates | Habitat and elevation |
|-----------|------------------------------|---------------------|---|
| A | Thathon | 20° 03'N, 99° 20'E | Lowland rice paddy and degraded marshland, 480 m |
| B | Doi Pha Hom Pok | 20° 05'N, 99° 10'E | Hill evergreen forest and secondary growth, 1400–2200 m |
| C | Fang | 19° 55'N, 99° 14'E | Lowland rice paddy, 480 m |
| D | Chiang Mai University Campus | 18° 47'N, 98° 58'E | Swampy scrub and grassland, 300 m |
| E | Doi Inthanon | 18° 35'N, 98° 29'E | Hill evergreen forest, 1650–2590 m Secondary growth and scrub, 1300 m. Deciduous dipterocarp forest and bamboo, 825 m |
| F | Beung Boraphet | 15° 43'N, 100° 14'E | Freshwater swamp and scrub, 30 m |
| G | Khao Yai | 14° 26'N, 101° 22'E | Dry evergreen forest, 800 m |
| H | Bangpoo | 13° 30'N, 100° 45'E | Coastal mangrove; freshwater marsh, 0–3 m |
| J | Samut Sakhon | 13° 29'N, 100° 15'E | Coastal mangrove, 0–3 m |
| K | Bang Phra | 13° 12'N, 101° 01'E | Scrub and grassland adjacent to freshwater lake, 50 m |
| L | Khao Sam Roi Yot | 12° 08'N, 99° 55'E | Freshwater marsh, 0–10 m |

All birds were weighed using Pesola spring balances, birds heavier than 50 g to the nearest 1 g (occasionally to the nearest 0.5 g) while those lighter than this were usually weighed to the nearest 0.1 g. Specimens retained for preparation as museum skins were sexed by dissection and the gonads measured *in situ* using Vernier callipers. Testes were noted as small (inactive) or enlarged (active) and the length of the larger testis was usually measured. Ovaries were noted as inactive unless any individual ova exceeded 0.5 mm in diameter, when the largest ovum was measured. Sex was not recorded for birds released, even in those species with sexually dimorphic plumages, as plumage characters may not always be wholly reliable (e.g. a female *Niltava grandis* had blue 'male' plumage on the head and one of the rectrices was blue on the outer web).

Weights

The weight data are too few to permit comparisons between sexes or age classes or to relate to breeding condition or to time of year, although in a few migratory species (e.g. *Phylloscopus fuscatus*), there is a suggestion of pre-migratory weight increase in the May samples (Table 2). Interpretation of weights from mixed samples of live and dead birds may need some care; von Bröckel (1973) found that a sample of Garden Warblers *Sylvia borin* weighed within 15 minutes of death averaged more than 1 g (5.5%) lighter than live birds.

Data on weight in relation to time of capture are available for 36 *Alcippe castaneiceps* and 49 *A. morrisonia*, 2 mainly insectivorous, forest-living species, caught between 09:00 and 18:00 (data pooled for sites B and E and the months January, February, April and December). Both species showed a gradual increase in mean weights throughout the day (*A. castaneiceps* $F_{1,34} = 4.44$, $p < 0.05$; *A. morrisonia* $F_{1,47} = 6.36$, $p < 0.025$; see Figs 1 and 2 — see Sokal & Rohlf 1981: 471). Although Nisbet & Medway (1972) found that *Acrocephalus (arundinaceus) orientalis* wintering in reedbeds in lowland Malaysia appeared to be most active in



the first 2–3 hours after sunrise, they did not find any marked changes in mean weights throughout the day. There appears to be no other published information on diurnal weight changes in birds in Southeast Asia.

Gonads

Deignan (1945) noted that the number of species breeding in North-West Thailand increased from November to a peak in April and May, this observation being based primarily on the condition of gonads and the collection of juveniles. Round (1982), working in the same area, found that the onset of nesting for 78 mainly insectivorous species inhabiting hill forest was in February and the relevant gonad data presented here support this conclusion. For many of the forest birds reported both here and by Deignan (1945), enlarged gonads were first recorded in April by Deignan, but were found earlier during this study. Gonad condition was not always recorded by Deignan's collectors and it seems possible that insufficient data resulted in him recording a later start to breeding.

Round (1982) discussed the timing of breeding in NW Thailand and noted that it might be advantageous for hill birds to rear their young to virtual independence before the onset of the rains in May, even though a peak in insect abundance may not occur until later. In moderate elevation forest (600–700 m) at Khao Yai, Central Thailand, McClure (1974b) found peak nesting activity among insectivorous birds between March and June. Herbert (1923–26) found that most birds in the lowlands around Bangkok had nests and eggs during the first half of the wet season, from May to July.

TABLE 2

(R) = resident; (M) = migrant species or subspecies. For key to sites, see Table 1. Data on weights and gonads correspond in serial order. For larger samples, only the range of weights is given. For gonads + = active; - = inactive. Small = testis length < 1.0 mm. In females: 8.3/3.0, for example, indicates ovary 8.3 mm and diameter of largest ovum 3.0 mm.

| Species | Site | Month | Sex | n | Weights (g) | | | Gonads activity | Gonads length (mm) |
|------------------------------------|------|-------|-----|---------------------------------------|--|------------|----|--------------------|---------------------------------|
| | | | | | | mean | SD | | |
| <i>Butorides striatus</i> (R) | J | 2 | — | 5 | 193.5, 209.5, 214.5, 226.5, 232.5 | 200 ± 24 | | | |
| | H | 9 | — | 1 | 170 | | | | |
| | H | 11 | — | 1 | 180.5 | | | | |
| | J | 11 | — | 1 | 173 | | | | |
| <i>Ixobrychus sinensis</i> (R) | F | 2 | M | 1 | 81 | | | — | 3.0 |
| <i>Ixobrychus cinnamomeus</i> (R) | F | 1 | — | 1 | 89.5 | | | | |
| | F | 2 | — | 1 | 131 | | | | |
| <i>Arborophila rufogularis</i> (R) | B | 1 | M | 1 | 212 | | | — | 6.0 |
| <i>Turnix tanki</i> (R) | B | 4 | F | 1 | 85 | | | + | 8.3/3.0 |
| <i>Treron sphenura</i> (R) | E | 12 | — | 1 | 280 | | | | |
| <i>Ducula badia</i> (R) | E | 12 | M | 1 | 510 | | | — | 14.0 |
| <i>Chalcophaps indica</i> (R) | B | 1 | F | 1 | 110 | | | + | 12.0/2.0 |
| <i>Cacomantis merulinus</i> (R) | D | 12 | — | 1 | 27 | | | | |
| <i>Caprimulgus affinis</i> (R) | E | 5 | F | 1 | 89 | | | | 7.7 |
| <i>Halcyon smyrnensis</i> (R) | D | 12 | — | 1 | 86 | | | | |
| <i>Halcyon pileata</i> (M) | J | 2 | — | 1 | 89 | | | | |
| | H | 10 | — | 2 | 83, 84.5 | | | | |
| | H | 11 | — | 1 | 92 | | | | |
| <i>Halcyon chloris</i> (R) | J | 2 | — | 1 | 67 | 62 ± 5 | | | |
| | J | 3 | — | 1 | 59.5 | | | | |
| | H | 9 | — | 3 | 62, 65, 65 | | | | |
| | H | 10 | — | 1 | 65 | | | | |
| | H | 11 | — | 2 | 51, 58 | | | | |
| <i>Merops philippinus</i> (R) | L | 5 | F | 1 | 32.8 | | | + | 11.5/5.4 |
| <i>Merops orientalis</i> (R) | D | 12 | — | 1 | 17.3 | | | | |
| <i>Megalaima franklinii</i> (R) | B | 1 | M | 1 | 81 | | | + | 3.5 |
| | B | 2 | F | 1 | 85.5 | | | + | 10.2/2.1 |
| | E | 12 | F | 2 | 76, 93 | | | — | 9.8, 8.0 |
| <i>Megalaima asiatica</i> (R) | E | 12 | F | 1 | 78.0 | | | — | 7.3 |
| <i>Jynx torquilla</i> (M) | D | 12 | — | 1 | 37.3 | | | | |
| | K | 12 | — | 1 | 32.4 | | | | |
| <i>Picumnus innominatus</i> (R) | B | 4 | F | 1 | 9.5 | | | + | 3.4/0.5 |
| | B | 4 | — | 1 | 8.6 | | | | |
| <i>Sasia ochracea</i> (R) | B | 4 | M | 1 | 8.9 | | | + | 3.5 |
| B | 1 | F | 1 | 9.9 | | | | — | 2.5 |
| B | 4 | F | 2 | 9.1, 10.5 | | | | + | 3.2/-, 4.7/2.2 |
| <i>Riparia riparia</i> (M) | F | 3 | M | 1 | 12.4 | | | — | |
| | F | 2 | — | 2 | 9.5, 10.6 | | | | (small) |
| F | 3 | — | 4 | 11.5, 11.8, 12.4, 13.0 | | | | | |
| <i>Hirundo rustica</i> (M) | F | 2 | M | 3 | 12.8, 13.8, 14.7 | 14.2 ± 1.0 | 3 | — | 1.5, (small), 3.0 |
| | F | 3 | M | 13 | Range 12.7 to 16.6 | | 13 | — | Range, small to 2.0 |
| F | 2 | F | 6 | 11.8, 12.3, 12.5, 13.1, 13.5, 13.6 | 12.8 ± 0.7 | 6 | | | 3.5, 4.0, 4.5, 5.0, 6.0, 4.0 |
| | F | 3 | F | 9 | 12.2, 12.9, 13.0, 13.5, 13.7, 14.2, 14.4, 14.4, 14.5 | 9 | | (small) | |
| F | 1 | — | 78 | Range 11.3 to 15.2 | 13.3 ± 0.8 | | | | |
| F | 2 | — | 248 | Range 11.5 to 16.4 | 13.6 ± 0.9 | | | | |
| F | 3 | — | 143 | Range 12.0 to 16.6 | 13.9 ± 1.0 | | | | |

| Species | Site | Month | Sex | n | Weights (g) | | | mean | SD | n | Gonads activity | length (mm) |
|--------------------------------------|------|-------|-----|----|--|--|--|------------|----|---|-----------------|--|
| | | | | | | | | | | | | |
| <i>Coracina polioptera</i> (R) | B | 1 | — | 1 | 35.8 | | | | | | | |
| <i>Pericrocotus solaris</i> (R) | B | 1 | — | 1 | 14.6 | | | | | | | |
| <i>Pericrocotus brevirostris</i> (R) | B | 4 | F | 1 | 19.8 | | | | | | | |
| <i>Aegithina tiphia</i> (R) | H | 4 | M | 1 | 12.3 | | | | | | | |
| | H | 4 | F | 1 | 12.0 | | | | | | | |
| | H | 4 | — | 1 | 13.4 | | | | | | | |
| | H | 9 | — | 1 | 13.6 | | | | | | | |
| <i>Pycnonotus striatus</i> (R) | E | 12 | M | 3 | 44.0, 45.5, 47.8 | | | | | 4 | — | 2.8, 4.6, 2.2, 1.9 |
| | E | 12 | F | 5 | 40.5, 42.9, 44.3, 45.0, 45.0 | | | 43.5 ± 1.9 | | 5 | — | 8.0, 6.5, 5.1, 6.4, 4.6 |
| | E | 12 | — | 6 | 43.0, 45.5, 45.5, 46.0, 47.0, 48.5 | | | 45.9 ± 1.8 | | | | |
| <i>Pycnonotus melanicterus</i> (R) | E | 1 | M | 1 | 28.4 | | | | | 1 | — | (small) |
| | E | 12 | M | 1 | 30.8 | | | | | 1 | — | (small) |
| | B | 2 | F | — | — | | | | | 1 | — | 4.7 |
| <i>Pycnonotus jocosus</i> (R) | B | 4 | M | 2 | 26.3, 27.6 | | | | | 2 | + | 7.3, 7.5 |
| | B | 4 | F | 8 | 23.3, 24.2, 24.5, 24.9, 25.0, 25.7, 27.6, 27.7 | | | 25.4 ± 1.6 | | 8 | + | 4.3/2.0, 6.1/1.7, 5.0/—, 5.5/2.0, 5.8/—, 7.0/3.0, 7.8/—, 7.9/1.8 |
| | B | 4 | — | 12 | Range 22.9 to 27.8 | | | 25.4 ± 1.4 | | | | |
| | D | 12 | — | 4 | 25.6, 27.9, 28.8, 29.2 | | | | | | | |
| <i>Pycnonotus xanthorrhous</i> (R) | B | 4 | M | 5 | 26.2, 26.3, 26.3, 28.9, 31.1 | | | 27.8 ± 2.2 | | 5 | + | 7.9, 7.3, 6.0, 8.1, 7.8 |
| | B | 4 | F | 7 | 24.4, 24.9, 25.1, 26.0, 27.5, 27.9, 28.5 | | | 26.3 ± 1.6 | | 7 | + | —/4.5, 5.7/1.3, 6.0/1.8, 5.8/—, 6.0/2.0, 10.4/3.1, 6.8/3.0 |
| <i>Pycnonotus aurigaster</i> (R) | B | 4 | M | 1 | 41.8 | | | | | 1 | + | 10.7 |
| | E | 1 | F | 1 | 29.7 | | | | | 1 | — | 6.4 |
| | B | 4 | F | 1 | 32.5 | | | | | 1 | — | 5.8 |
| <i>Pycnonotus flavescens</i> (R) | B | 4 | M | 2 | 28.9, 28.9 | | | | | 2 | + | 6.0, 6.1 |
| | E | 12 | M | 1 | 35.0 | | | | | 1 | — | 2.7 |
| | B | 4 | F | 6 | 27.1, 27.8, 27.9, 28.0, 28.1, 29.4 | | | 28.9 ± 2.1 | | 6 | + | 6.9/2.4, —/1.8, 6.9/2.2, 7.6/2.4, 8.4/2.1, 9.9/3.0 |
| | E | 12 | F | 1 | 31.6 | | | | | 1 | — | |
| | B | 4 | — | 3 | 27.6, 28.0, 28.0 | | | | | | | 6.5 |
| <i>Pycnonotus goiavier</i> (R) | K | 9 | — | 1 | 28.1 | | | | | | | |
| <i>Pycnonotus blanfordi</i> (R) | F | 2 | — | 1 | 32.1 | | | | | | | |
| | F | 3 | — | 1 | 33.4 | | | | | | | |
| | D | 12 | — | 6 | 31.3, 33.2, 33.4, 33.9, 34.0, 35.7 | | | 33.4 ± 1.3 | | | | |
| <i>Criniger pallidus</i> (R) | G | 11 | — | 10 | 41, 43, 44, 45, 46, 47, 48, 49, 52 | | | 46 ± 3 | | | | |
| <i>Hypsipetes propinquus</i> (R) | E | 1 | F | 1 | 25.9 | | | | | 1 | — | 5.9 |
| <i>Hypsipetes mcclellandii</i> (R) | B | 2 | M | 1 | 36.6 | | | | | 1 | + | 7.0 |
| | B | 4 | M | 4 | 31.9, 34.6, 37.5, 37.7 | | | 35.4 ± 2.5 | | 4 | + | 8.0, 8.0, 8.5, 9.5 |
| | E | 12 | M | 2 | 32.2, 37.2 | | | | | 2 | — | 2.0, 3.0 |
| | B | 1 | F | 2 | 32.5, 36.3 | | | | | 2 | + | 9.0/1.0, 7.0/1.8 |
| | B | 2 | F | 2 | 36.2, 38.2 | | | 35.1 ± 2.4 | | 2 | + | 9.5/5.0, 9.5/2.0 |
| | B | 4 | F | 1 | 34.7 | | | | | 1 | + | 8.0/2.0 |
| | E | 12 | F | 3 | 31.3, 36.5 | | | | | 2 | — | (small), 5.5 |
| | E | 12 | — | 3 | 37.4, 42.7, 43.5 | | | | | | | |
| <i>Hypsipetes thompsoni</i> (R) | E | 12 | M | 1 | 41.8 | | | | | 1 | — | (small) |
| <i>Dicrurus remifer</i> (R) | B | 4 | M | — | 45.4 | | | | | 1 | + | 10.3 |
| | B | 4 | — | 1 | 46.0 | | | | | | | |
| <i>Dendrocitta formosae</i> (R) | B | 4 | F | 1 | 80 | | | | | 1 | + | 7.5/1.0 |
| <i>Parus spilonotus</i> (R) | B | 4 | M | 1 | 16.7 | | | | | 1 | — | 1.0 |
| | E | 12 | M | 1 | 17.0 | | | | | 1 | + | 3.0 |
| | E | 12 | F | 1 | 15.8 | | | | | 1 | — | 5.0 |
| <i>Sylviparus modestus</i> (R) | B | 2 | F | 1 | 7.1 | | | | | 1 | + | —/4.4 |
| | E | 12 | F | 2 | 6.2, 6.5 | | | | | 2 | — | (small) |
| <i>Sitta nagaensis</i> (R) | B | 1 | M | — | 13.9 | | | | | 1 | + | 4.5 |
| | B | 4 | M | 1 | 15.5 | | | | | 1 | + | 4.4 |
| <i>Certhia discolor</i> (R) | B | 4 | M | 1 | 9.5 | | | | | 1 | — | (small) |
| | B | 4 | F | 1 | 9.3 | | | | | 1 | — | |
| | E | 12 | F | 1 | 9.7 | | | 9.7 ± 0.3 | | 1 | — | 3.3 |
| | B | 1 | — | 2 | 9.5, 10.2 | | | | | | | 6.0 |
| | B | 4 | — | 1 | 9.8 | | | | | | | |
| | E | 12 | — | 1 | 9.8 | | | | | | | |
| <i>Pellorneum ruficeps</i> (R) | E | 1 | M | 1 | 23.6 | | | | | 1 | — | (small) |
| | E | 1 | F | 1 | 20.6 | | | | | 1 | — | 5.1 |

| Species | Site | Month | Sex | n | Weights (g) | | | n | Gonads activity | length (mm) | |
|--------------------------------------|------|-------|-----|----|--|----|--|------------|-----------------|--|--|
| | | | | | mean | SD | | | | | |
| <i>Pellorneum albiventre</i> (R) | B | 4 | F | 1 | 18.4 | | | 1 | + | 6.8/1.0 | |
| <i>Trichastoma tickelli</i> (R) | B | 1 | M | 2 | 17.4, 17.7 | | | 2 | + | 2.0, 3.6 | |
| | B | 1 | F | 1 | 16.2 | | | 1 | + | 5.2/- | |
| <i>Pomatorhinus erythrogenys</i> (R) | B | 1 | M | 1 | 57.0 | | | 1 | + | 5.5 | |
| | E | 12 | F | 2 | 52.5, 56 | | | 2 | + | 10.0/5.0, 6.2/- | |
| | B | 4 | — | 1 | 54 | | | | | | |
| <i>Pomatorhinus schisticeps</i> (R) | B | 1 | M | 2 | 34.6, 36.2 | | | 2 | + | 4.0, 4.5 | |
| | E | 12 | M | 1 | 39.5 | | | 1 | - | 3.4 | |
| | B | 1 | F | 2 | 31.5, 35.2 | | | 2 | + | 8.0/1.5, 10.5/4.0 | |
| | B | 4 | F | 1 | 39.6 | | | 1 | + | 7.2/2.5 | |
| | E | 12 | F | 2 | 33, 40 | | | 2 | - | (small), 8.1 | |
| | B | 4 | — | 1 | 32.8 | | | | | | |
| <i>Napothera epilepidota</i> (R) | B | 1 | M | 2 | 15.4, 19.0 | | | 2 | - | (small), 1.8 | |
| | B | 1 | F | 3 | 14.5, 15.4, 15.5 | | | 3 | - | 5.2, 3.7, 3.7 | |
| <i>Pnoepyga pusilla</i> (R) | B | 2 | F | 1 | 12.9 | | | 1 | - | 5.5 | |
| | B | 4 | F | 1 | 12.5 | | | 1 | + | 5.2/1.0 | |
| <i>Stachyris rufifrons</i> (R) | B | 1 | M | 2 | 8.5, 8.7 | | | 2 | + | 3.5, 7.0 | |
| | E | 1 | M | 1 | 8.5 | | | 1 | - | 5.2 | |
| | B | 1 | F | 1 | 7.5 | | | 1 | + | 6.1/1.5 | |
| | E | 1 | F | 1 | 7.5 | | | 1 | - | 4.2 | |
| <i>Stachyris chrysaea</i> (R) | B | 1 | M | 2 | 7.8 ^a , 8.0 | | | 2 | + | 1.0 ^a (small, juv), 6.5 | |
| | B | 4 | M | 2 | 7.8, 8.0 | | | 1 | + | | |
| | E | 12 | M | 2 | 7.5, 8.1 | | | 2 | - | 5.0, 1.5 | |
| | B | 1 | F | 4 | 7.0, 7.4, 8.5, 11.0 | | | 4 | + | 4.7/-, 4.0/1.5, 6.0/1.5, -/4.0, | |
| | B | 4 | F | 1 | 7.3 | | | 1 | - | 5.0 | |
| | E | 12 | F | 2 | 7.6, 8.1 | | | 2 | - | 4.5, 5.0 | |
| <i>Stachyris nigriceps</i> (R) | B | 1 | M | 5 | 14.9, 16.2, 16.5, 16.5, 17.0 | | | 5 | + | 10.0, 10.5, 7.0, 8.5, 10.0 | |
| | B | 4 | M | 4 | 14.6, 15.3 ^a , 16.0 ^a , 16.1 | | | 4 | + | 9.7, 1.3 ^a , 1.3 ^a , 9.8 | |
| | E | 12 | M | 2 | 14.5, 16.0 | | | 2 | + | 2.5, 10.0 | |
| | B | 1 | F | 6 | 13.1, 14.6, 14.8, 15.0, 16.0, 19.0 | | | 6 | + | 5.0/-, 6.5/1.0, 5.5/-, 5.0/-, 4.5/-, 5.6/- | |
| | B | 4 | F | 2 | 11.9, 14.8 | | | 3 | + | 5.0/-, 5.9/1.8, 6.0/- | |
| | E | 12 | F | 3 | 14.1, 15.0, 15.7 | | | 2 | + | 5.0/1/-, 4.0/0, (n.m.) | |
| | E | 12 | — | 2 | 15.8, 16.3 | | | | | | |
| <i>Macronous gularis</i> (R) | E | 1 | M | 2 | 10.0, 10.7 | | | 2 | + | 5.2, 1.7 | |
| | E | 1 | F | 1 | 9.5 | | | 1 | - | 3.6 | |
| <i>Chrysomma sinense</i> (R) | F | 2 | M | 2 | 15.7, 16.5 | | | 2 | - | 1.7, 1.9 | |
| | F | 2 | F | 2 | 14.5, 15.6 | | | 2 | - | 4.0, 4.5 | |
| | E | 12 | F | 2 | 17.7, 21.0 | | | 1 | - | 5.8, (n.m.) | |
| | A | 2 | — | 3 | 17.6, 18.5, 19.5 | | | | | | |
| | F | 3 | — | 2 | 16.0, 16.4 | | | | | | |
| | K | 9 | — | 6 | 17.0, 17.3, 17.6, 17.8, 18.5, 20.1 | | | | | | |
| | K | 12 | — | 6 | 15.8, 16.0, 17.0, 17.6, 17.8, 20.1 | | | | | | |
| | D | 12 | — | 2 | 18.7, 21.5 | | | | | | |
| <i>Timalia pileata</i> (R) | A | 2 | — | 1 | 18.5 | | | | | | |
| | K | 9 | — | 1 | 21.5 | | | | | | |
| | D | 12 | — | 3 | 20.4, 20.8, 21.0 | | | 20.4 ± 1.2 | | | |
| <i>Garrulax erythrocephalus</i> (R) | B | 2 | M | 3 | 81.0, 83.5, 85.0 | | | 3 | - | 3.2, 2.5, 3.5 | |
| | B | 4 | M | 1 | 77 | | | 1 | + | 5.9 | |
| | E | 12 | M | 1 | 76.5 | | | 1 | - | (small) | |
| | B | 1 | F | 2 | 72.3, 75.0 | | | 2 | - | 8.0/1.0, 7.8/- | |
| | E | 12 | F | 2 | 70.0, 71.5 | | | 2 | - | (small), 7.5/- | |
| | E | 12 | — | 4 | 78.5, 79.0, 79.5, 89.5 | | | | | | |
| <i>Liocichla phoenicea</i> (R) | B | 1 | M | 1 | 47.0 | | | 1 | - | (small) | |
| | B | 4 | M | 1 | 47.0 | | | 1 | + | 10.4 | |
| | B | 1 | F | 1 | 41.6 | | | 1 | - | 6.3 | |
| | B | 2 | F | 1 | 47.9 | | | 1 | - | 5.1 | |
| | B | 4 | F | 1 | 44.5 | | | 1 | + | 8.8/1.5 | |
| | B | 4 | — | 1 | 55.0 | | | | | | |
| <i>Leiothrix argentauris</i> (R) | B | 4 | M | 3 | 23.8, 24.8, 25.6 | | | 3 | + | 11.5, 12.4, 8.5 | |
| | E | 12 | M | 2 | 25.5, 26.0 | | | 2 | - | (small), (small) | |
| | B | 4 | F | 2 | 26.4, 26.6 | | | 2 | + | -3.5, -/8.4 | |
| | E | 12 | F | 2 | 23.8, 24.5 | | | 2 | - | 5.8, 7.5 | |
| | E | 12 | — | 13 | 24.2, 24.3, 24.8, 25.3, 25.5, 25.6, 25.6, 25.7 26.2, 26.5, 26.7, 27.0, 27.5 | | | | | | |
| <i>Pteruthius flaviscapus</i> (R) | B | 2 | F | 1 | 32.6 | | | 1 | - | 6.3 | |
| <i>Actinodura ramsayi</i> (R) | B | 1 | M | 1 | 39.5 | | | 1 | + | 6.4 | |
| | B | 4 | M | 3 | 37.4, 37.9, 38.5 | | | 3 | + | 8.8, 9.6, 8.0 | |
| | B | 4 | F | 1 | 35.5 | | | 1 | - | 8.0 | |
| | E | 12 | F | 1 | 41.3 | | | 1 | - | 6.6 | |

| Species | Site | Month | Sex | n | Weights (g) | | mean | SD | n | Gonads | |
|------------------------------|------|-------|-----|----|---|--|------------|----|----|----------|---|
| | | | | | | | | | | activity | length (mm) |
| Minla cyanouroptera (R) | B | 2 | M | 1 | 15.4 | | | | 1 | - | 4.3 |
| | B | 4 | M | 1 | 17.3 | | | | 1 | + | 5.7 |
| | B | 1 | F | 1 | 15.8 | | | | 1 | - | 5.1 |
| | B | 2 | F | 1 | 15.8 | | | | 1 | + | 5.0/1.0 |
| Minla strigula (R) | E | 12 | M | 5 | 19.0, 19.5, 20.5 21.6, 21.9 | | 20.5 ± 1.3 | | 5 | - | (small), (small), 2.0, (small), 2.5 |
| | E | 12 | F | 7 | 18.5, 18.7, 18.7, 19.2, 19.2, 19.3, 20.3 | | 19.1 ± 0.6 | | 7 | - | (small), 3.0, 5.0, (small), 6.0, (small), (small) |
| | E | 5 | - | 8 | 19.0, 19.0, 19.2, 19.3, 20.5, 20.5, 20.8, 20.8 | | 19.9 ± 0.8 | | | | |
| | E | 12 | - | 10 | 18.2, 18.5, 18.7, 19.3, 19.3, 19.6, 19.8, 19.8, 19.9, 20.4 | | 19.4 ± 0.7 | | | | |
| Alcippe castaneiceps (R) | B | 1 | M | 3 | 10.0, 10.3, 10.7 | | | | 3 | + | 3.5, 5.0, 2.5 |
| | B | 2 | M | 14 | 9.2, 9.2, 9.5, 9.5 9.6, 9.7, 9.9, 9.9, 10.4, 10.5, 10.5, 10.6 10.8, 11.2 | | 10.1 ± 0.6 | | 14 | + | 2.5, 2.4, 6.5, 2.0, 3.0, 4.0, 6.0, 6.5, 6.4, 7.5, 6.5, 6.0, 2.5, 6.0 |
| | B | 4 | M | 1 | 9.4 | | | | 1 | + | 5.8 |
| | E | 12 | M | 14 | 9.0, 9.2, 9.4, 9.5 9.6, 9.7, 10.0, 10.4 10.4, 10.6, 10.8, 10.8, 11.2, 11.2 | | 10.1 ± 0.7 | | 14 | - | 1.5, 1.5, 2.0, (small), 1.5, 2.5, 1.0, 2.0, (small), (small), 1.5, 2.0, 2.5, (small) |
| | B | 1 | F | 1 | 11.5 | | | | 1 | + | 18.0/4.5 |
| | B | 2 | F | 6 | 8.9, 9.2, 9.5, 9.7, 9.8, 10.0 | | 9.8 ± 0.8 | | 6 | + | 4.5/1.0, 5.0/1.0, 5.0/1.0, 4.7/1.3, 4.0/-, 6.0/1.7 |
| | E | 12 | F | 10 | 9.2, 9.2, 9.5, 9.7, 10.0, 10.2, 10.3, 10.4, 10.5, 10.5 | | 10.0 ± 0.5 | | 10 | - | 4.5, 4.0, (small), 5.0, 4.5, 4.5, (small), (small), (small) |
| | B | 2 | - | 6 | 9.7, 9.7, 9.9, 10.0, 10.3, 10.7 | | 10.0 ± 0.7 | | | | |
| | B | 4 | - | 3 | 8.6, 10.5, 10.7 | | | | | | |
| | E | 12 | - | 23 | Range 8.3 to 10.5 | | 9.5 ± 0.6 | | | | |
| Alcippe poiocephala (R) | E | 1 | F | 1 | 15.4 | | | | 1 | - | 5.9 |
| Alcippe morrisonia (R) | B | 1 | M | 9 | 13.2, 13.7, 13.9, 14.5, 14.5, 15.1, 15.2, 15.5, 15.5 | | 14.7 ± 0.7 | | 9 | + | 2.5, 3.0, (small), 3.0, 5.0, 2.0, 6.0, 5.5, 2.0 |
| | B | 2 | M | 5 | 13.9, 14.8, 14.8, 14.9, 15.6 | | | | 5 | + | 6.0, 4.0, 6.0, 4.4, 5.1 |
| | B | 4 | M | 3 | 13.8, 14.0, 14.1 | | | | 3 | + | 6.0, 4.0, 7.7 |
| | E | 12 | M | 6 | 13.0, 13.6, 14.0, 14.2, 14.2, 15.1 | | 14.0 ± 0.7 | | 6 | - | 2.2, (small), 1.5, 2.0, 3.0, 2.5 |
| | B | 1 | F | 3 | 13.5, 14.6, 15.7 | | 14.5 ± 0.8 | | 3 | - | 5.5, 5.5, 4.5 |
| | B | 2 | F | 4 | 13.9, 13.9, 14.8, 14.9 | | | | 4 | + | 5.2/1.4, 5.3/1.4, 6.0/2.0, 6.5/1.0 |
| | B | 4 | F | 3 | 13.7, 14.6, 16.8 | | | | 3 | + | 5.0/1.0, 6.0/1.8, 8.0/2.0 |
| | E | 12 | F | 4 | 12.5, 12.8, 14.3, 14.4 | | | | 4 | - | 4.5, 4.5, 5.5, (small) |
| | B | 1 | - | 20 | Range 13.6 to 15.9 | | 14.5 ± 0.7 | | | | |
| | B | 4 | - | 10 | Range 13.4 to 15.5 | | 14.2 ± 0.5 | | | | |
| Heterophasia annectens (R) | E | 12 | - | 30 | Range 12.8 to 16.5 | | 14.8 ± 0.7 | | | | |
| | B | 4 | M | 1 | 24.2 | | | | 1 | + | 5.7 |
| | B | 1 | F | 1 | 24.8 | | | | 1 | + | 8.1/1.0 |
| Heterophasia melanoleuca (R) | B | 4 | M | 2 | 36.4, 36.4 | | | | 2 | + | 7.8, 9.0 |
| | E | 12 | M | 4 | 30.5, 31.1, 32.5, 33.0 | | 33.3 ± 2.6 | | 4 | - | 1.8, 1.7, 2.6, 2.5 |
| | B | 1 | F | 1 | 30.1 | | | | 1 | - | 7.0 |
| | B | 4 | F | 1 | 31.2 | | 31.7 ± 1.6 | | 1 | + | 9.5/2.0 |
| | E | 12 | F | 3 | 30.6, 32.8, 33.8 | | | | 3 | - | 4.1, 6.3, 7.4 |
| Yuhina flavicollis (R) | B | 2 | M | 2 | 15.2, 16.8 | | | | 2 | - | 3.0, 4.4 |
| | B | 4 | M | 2 | 14.0, 14.6 | | | | 2 | + | 6.3, 6.6 |
| | B | 2 | F | 1 | 14.9 | | | | 1 | + | 5.8/1.3 |
| | B | 4 | F | 2 | 12.5, 16.7 | | | | 2 | + | -/1.7, -/7.5 |
| Yuhina zantholeuca (R) | G | 11 | - | 1 | 11.2 | | | | | | |
| Paradoxornis gularis (R) | B | 1 | - | 1 | 27.7 | | | | 1 | + | 7.8 |
| | B | 4 | M | 1 | 26.1 | | | | 1 | + | 11.0 |
| | B | 4 | F | 1 | 35.5 | | | | | | |
| Brachypteryx leucophrys (R) | B | 4 | M | 1 | 13.6 | | | | 1 | + | 4.3 |
| | B | 1 | F | 1 | 15.2 | | | | 1 | - | 4.5 |
| | B | 4 | F | 2 | 13.9, 15.5 | | | | 2 | + | 6.6/1.6, 7.0/1.3 |
| Brachypteryx montana (R) | E | 4 | M | 1 | 20.8 | | | | 1 | + | 7.0 |
| | E | 4 | - | 1 | 18.6 | | | | | | |

| Species | Site | Month | Sex | n | Weights (g) | Weights (g) | | | Gonads | length (mm) |
|-------------------------------------|------|-------|-----|----|--|-------------|------------|------------|----------|-------------------------------------|
| | | | | | | mean | SD | n | activity | |
| <i>Erithacus calliope</i> (M) | B | 4 | F | 1 | 22.2 | 20.1 ± 1.4 | 20.1 ± 1.4 | 1 | - | 4.5 |
| | F | 1 | - | 2 | 19.3, 20.5 | | | 2 | - | |
| | F | 2 | - | 5 | 18.8, 19.1, 19.5, 20.8, 21.1 | | | 1 | - | 1.5 |
| | F | 3 | - | 3 | 18.8, 19.8, 21.3 | | | 1 | - | 4.5 |
| | D | 12 | - | 3 | 18.0, 20.3, 23.4 | | | 2 | - | 4.2, 4.7 |
| | K | 12 | - | 3 | 18.8, 18.9, 21.0 | | | 2 | - | |
| | A | 2 | - | 18 | Range 13.6 to 18.0 | | | 15.2 ± 1.1 | | |
| | F | 2 | - | 2 | 15.3, 15.4 | | | 15.5 ± 0.4 | | |
| <i>Erithacus svecicus</i> (M) | F | 3 | - | 5 | 15.0, 15.2, 15.8, 15.8, 15.9 | | | 15.5 ± 0.4 | | |
| | F | 12 | - | 3 | 14.4, 15.3, 17.3 | | | | | |
| | E | 1 | M | 1 | 15.0 | 15.5 ± 0.9 | 15.5 ± 0.9 | 1 | - | (small) |
| | H | 9 | - | 1 | 14.1 | | | 2 | - | (both small) |
| | E | 12 | M | 1 | 12.5 | | | 1 | - | (small) |
| | B | 2 | F | 2 | 13.2, 14.7 | | | 2 | - | 4.5, 4.0 |
| | E | 12 | F | 1 | 13.9 | | | 2 | - | (both small) |
| | B | 2 | - | 1 | 12.4 | | | | | |
| <i>Copyschus saularis</i> (R) | H | 4 | - | 2 | 35.1, 38.0 | 26.5 ± 3.4 | 26.5 ± 3.4 | 2 | - | 2.0, (small), (small), 2.0, 1.5 |
| | D | 12 | - | 2 | 37.8, 40.0 | | | 2 | - | |
| <i>Cinclidium leucurum</i> (R) | B | 1 | M | 5 | 22.7, 25.5, 25.8, 26.5, 31.9 | 24.9 ± 0.7 | 24.9 ± 0.7 | 5 | - | 4.5, 3.8, 5.5, 3.5, 6.4 |
| | B | 4 | M | 5 | 24.3, 24.4, 24.6, 25.4, 25.8 | | | 5 | + | |
| | E | 12 | M | 2 | 20.5, 27.5 | | | 2 | - | (both small) |
| | E | 12 | - | 1 | 25.5 | | | | | |
| | F | 1 | - | 1 | 12.6 | | | | | |
| | F | 2 | - | 2 | 12.2, 12.5 | | | | | |
| | A | 2 | - | 9 | 11.8, 12.2, 12.8, 12.8, 12.8, 13.0, 13.3, 13.5, 13.6 | | | 12.8 ± 0.5 | | |
| | | | | | | | | | | |
| <i>Saxicola caprata</i> (R) | A | 2 | - | 2 | 13.0, 13.8 | | | | | |
| <i>Saxicola ferrea</i> (R) | E | 12 | M | 2 | 15.0, 15.7 | | | 2 | - | 2.0, (small) |
| <i>Myiohoneus caeruleus</i> (R) | E | 12 | F | 1 | 13.4 | | | 1 | - | 5.5 |
| | B | 2 | - | 1 | 12.7 | | | | | |
| <i>Zoothera marginata</i> (R) | E | 12 | F | 1 | 8.1 | | | 1 | - | 7.5 |
| <i>Turdus obscurus</i> (M) | B | 4 | M | 1 | 5.5 | | | 1 | - | 2.8 |
| | E | 4 | F | 1 | 6.7 | | | 1 | - | 7.0/0.5 |
| <i>Gerygone sulphurea</i> (R) | H | 4 | F | 2 | 5.5, 5.8 | | | 2 | + | 3.3/0.5, 4.5/0.5 |
| | B | 1 | M | 3 | 6.9, 6.9, 7.0 | | | 3 | - | (small), 1.0, 0.9 |
| | B | 4 | M | 1 | 7.6 | | | 1 | - | 2.0 |
| | E | 12 | M | 2 | 6.8, 7.0 | | | 2 | - | (both small) |
| | B | 4 | F | 2 | 6.3, 6.6 | | | 2 | - | 3.5, 2.1 |
| | B | 1 | - | 2 | 6.3, 8.0 | | | | | |
| | E | 12 | - | 4 | 6.2, 6.9, 7.2, 7.5 | | | | | |
| | | | | | | | | | | |
| <i>Abroscopus superciliaris</i> (R) | E | 1 | M | 1 | 5.8 | | | 1 | - | (small) |
| <i>Phylloscopus subaffinis</i> (M) | E | 12 | M | 1 | 6.2 | | | 1 | - | (small) |
| <i>Phylloscopus fuscatus</i> (M) | H | 4 | F | 2 | 7.0, 8.8 | | | 2 | - | 3.0, 3.5 |
| | F | 1 | - | 3 | 7.1, 8.3, 9.4 | | | | | |
| | A, F | 2 | - | 18 | Range 7.0 to 9.4 | | | 7.9 ± 0.6 | | |
| | H | 4 | - | 1 | 10.8 | | | | | |
| | D | 12 | - | 2 | 7.5, 9.4 | | | | | |
| | K | 12 | - | 2 | 8.6, 9.4 | | | | | |
| | | | | | | | | | | |
| <i>Phylloscopus armandii</i> (M) | B | 4 | M | 1 | 9.6 | | | 1 | - | 1.5 |
| | E | 12 | M | 1 | 9.4 | | | 1 | - | (small) |
| | E | 12 | F | 1 | 9.0 | | | 1 | - | 3.8 |
| <i>Phylloscopus schwarzi</i> (M) | B | 4 | F | 1 | 9.3 | | | 1 | - | 9.0 |
| | B | 2 | M | 7 | 6.3, 6.3, 6.3, 6.4 | | | | | |
| <i>Phylloscopus pulcher</i> (M) | | | | | 6.4, 6.4, 6.6 | | | 6.2 ± 0.4 | | 0.5, 1.0, (small), 0.5, |
| | E | 12 | M | 2 | 5.5, 5.5 | | | 2 | - | 0.5, (small), (small), (both small) |
| | B | 1 | F | 1 | 5.9 | | | 1 | - | 3.3 |
| | E | 12 | - | 1 | 5.6 | | | 3 | - | (all small) |
| | E | 12 | - | 1 | 5.5 | | | | | |
| | B | 1 | - | 1 | 5.5 | | | | | |
| | D | 12 | - | 3 | 5.3, 5.4, 5.5 | | | | | |
| | | | | | | | | | | |
| <i>Phylloscopus inornatus</i> (M) | E | 12 | M | 3 | 4.8, 6.0, 6.0 | | | | | |
| | E | 12 | F | 2 | 5.0, 5.4 | | | | | |
| | E | 12 | - | 1 | 5.6 | | | | | |
| | B | 1 | - | 1 | 5.5 | | | | | |
| | D | 12 | - | 3 | 5.3, 5.4, 5.5 | | | | | |

| Species | Site | Month | Sex | n | Weights (g) | Weights (g) | | | Gonads | length (mm) | |
|--------------------------------------|------|-------|-----|----|---|-------------|----|-----|--|------------------------------------|--|
| | | | | | | mean | SD | n | activity | | |
| <i>Phylloscopus proregulus</i> (M) | B | 1 | M | 1 | 5.2 | 5.4 ± 0.3 | 1 | — | (small) | 2.3 3.0 | |
| | B | 4 | M | 1 | 5.8 | | | 1 | + | | |
| | B | 1 | F | 1 | 5.6 | | | 1 | — | | |
| | B | 4 | F | 1 | 5.4 | | | 2 | — | | |
| | B | 4 | — | 1 | 5.1 | | | 1 | — | | |
| <i>Phylloscopus maculipennis</i> (R) | E | 5 | M | 3 | 4.3, 4.8, 4.9 | 4.7 ± 0.2 | 2 | — | (n.m.) | 2.0, 2.5 4.0 | |
| | E | 12 | M | 2 | 4.7, 4.8 | | | 1 | — | | |
| | E | 12 | F | 1 | 4.6 | | | 2 | — | | |
| | E | 5 | — | 3 | 4.5, 4.6, 5.1 | | | 1 | — | | |
| | E | 12 | — | 4 | 4.5, 4.8, 4.8, 4.8 | | | 2 | — | | |
| <i>Phylloscopus borealis</i> (M) | H | 9 | — | 1 | 10.1 | | | | | | |
| <i>Phylloscopus trochiloides</i> (M) | E | 12 | F | 7 | 7.3 | | | 1 | — | (small) | |
| <i>Phylloscopus reguloides</i> (M) | B | 1 | M | 1 | 8.5 | | | 1 | — | (small) | |
| | B | 2 | — | 1 | 6.9 | | | | | | |
| <i>Phylloscopus davisoni</i> (R) | B | 1 | M | 2 | 6.3, 8.1 | 6.4 ± 0.7 | 2 | + | 3.0, 1.5 3.1 2.9 3.3 5.7, 2.6 | | |
| | B | 4 | M | 1 | 6.3 | | | 1 | + | | |
| | E | 12 | M | 1 | 6.4 | | | 1 | — | | |
| | B | 4 | F | 1 | 6.8 | | | 1 | — | | |
| | E | 12 | F | 2 | 5.8, 6.5 | | | 2 | — | | |
| | B | 1 | — | 1 | 6.2 | | | 1 | — | | |
| | B | 4 | — | 1 | 5.7 | | | 1 | — | | |
| <i>Acrocephalus aedon</i> (M) | D | 12 | — | 4 | 21.7, 22.6, 22.9, 23.2 | | | | | | |
| <i>Acrocephalus arundinaceus</i> (M) | F | 2 | M | 3 | 23.9, 27.2, 29.5 | 23.6 ± 3.6 | 2 | — | 1.5, (small), (small) | 2.3 6.5, 5.0 5.5 5.6, 6.5 | |
| | L | 5 | M | 2 | 28.5 | | | 1 | — | | |
| | F | 2 | F | 2 | 20.8, 21.1 | | | 2 | — | | |
| | H | 4 | F | 1 | 20.1 | | | 1 | — | | |
| | L | 5 | F | 2 | 18.4, 22.2 | | | 2 | — | | |
| | A | 2 | — | 2 | 23.9, 25.7 | | | | | | |
| | F | 3 | — | 2 | 20.5, 21.9 | | | | | | |
| | H | 4 | — | 2 | 23.0, 23.8 | | | | | | |
| | L | 5 | — | 25 | Range 18.3 to 31.3 | | | | | | |
| | K | 9 | — | 13 | Range 21.6 to 30.7 | | | | | | |
| <i>Acrocephalus bistrigiceps</i> (M) | K | 12 | — | 52 | Range 20.4 to 28.3 | 24.0 ± 1.7 | 3 | — | (all small) 1.0, 1.0, 1.0, 1.3, 1.0, 1.0, (small), (small), (small) | 1.0 3.7 4.8 | |
| | F | 1 | M | 3 | 7.8, 9.1, 9.4 | | | 9 | — | | |
| | F | 2 | M | 9 | 6.9, 7.1, 7.2, 7.7, 7.9, 8.3, 8.3, 8.3, 8.5 | | | | | | |
| | L | 5 | M | 1 | 7.3 | | | 1 | — | | |
| | F | 2 | F | 1 | 7.8 | | | 1 | — | | |
| | L | 5 | F | 1 | 7.6 | | | 1 | — | | |
| | A | 2 | — | 3 | 7.3, 8.0, 8.6 | | | | | | |
| | F | 1 | — | 1 | 11.6 | | | | | | |
| | F | 2 | — | 1 | 7.4 | | | | | | |
| | F | 3 | — | 1 | 7.2 | | | | | | |
| <i>Acrocephalus agricola</i> (M) | H | 4 | — | 5 | 7.6, 8.0, 8.0, 8.5, 8.9 | 8.2 ± 0.9 | 3 | — | 2.8 | | |
| | L | 5 | — | 8 | 7.6, 8.1, 8.3, 8.3, 8.6, 8.8, 8.9, 9.2, | | | 9 | — | | |
| | D | 12 | — | 1 | 9.3 | | | | | | |
| | K | 12 | — | 7 | 7.0, 7.2, 7.5, 8.1, 8.3, 8.6, 10.5 | | | | | | |
| | L | 5 | F | 1 | 7.9 | | | 1 | — | | |
| | A | 2 | — | 1 | 7.7 | | | | | | |
| | A | 2 | — | 1 | 7.7 | | | | | | |
| <i>Acrocephalus concinens</i> (M) | F | 2 | M | 8 | 8.9 | 14.0 ± 1.0 | 1 | — | 2.0 | | |
| | F | 3 | M | 1 | 8.7 | | | 1 | — | | |
| | F | 2 | F | 1 | 7.9 | | | 1 | — | | |
| | A | 2 | — | 1 | 7.7 | | | | | | |
| <i>Locustella certhiola</i> (M) | F | 1 | M | 1 | 12.1 | 14.0 ± 1.0 | 1 | — | (small) | 2.0 2.0 2.0 4.0 | |
| | F | 2 | M | 1 | 13.9 | | | 1 | — | | |
| | L | 5 | M | 1 | 14.9 | | | 1 | — | | |
| | F | 1 | F | 1 | 13.9 | | | 1 | — | | |
| | H | 4 | — | 1 | 14.7 | | | | | | |
| <i>Locustella lanceolata</i> (M) | F | 2 | F | 1 | 10.6 | 7.2 ± 0.7 | 1 | — | 3.6 | | |
| | A | 2 | — | 2 | 36.5, 39.0 | | | 1 | + | | |
| <i>Orthotomus sutorius</i> (R) | H | 4 | M | 1 | 7.0 | 7.2 ± 0.7 | 2 | + | 3.7 4.5, 8.0/2.0 | | |
| | H | 4 | F | 2 | 6.1, 6.8 | | | 1 | + | | |
| | H | 4 | — | 5 | 6.7, 7.0, 7.9, 8.0, 8.8 | | | 2 | + | | |
| | H | 9 | — | 4 | 6.6, 7.0, 7.1, 7.3 | | | | | | |
| | K | 9 | — | 1 | 7.4 | | | | | | |
| <i>Orthotomus cucullatus</i> (R) | B | 1 | M | 1 | 6.1 | 7.2 ± 0.7 | 1 | (+) | 2.0 | | |
| | B | 1 | F | 1 | 5.8 | | | 1 | — | | |
| | B | 1 | — | 2 | 5.3, 5.4 | | | | | | |

| Species | Site | Month | Sex | n | Weights (g) | Gonads | | | | |
|--|----------------|-------|-----|---|------------------------|------------|----|-------------|------------------|---|
| | | | | | | mean | SD | n | activity | length (mm) |
| <i>Prinia hodgsonii</i> (R) | E | 12 | M | — | 5.5 | — | — | I | — | 1.5 4.8 |
| | E | 12 | F | 1 | | | | | | |
| <i>Prinia subflava</i> (R) | H | 4 | M | 3 | 9.6, 10.1, 10.1 | 8.8 ± 1.4 | — | 3 1 | + — | 5.0, 5.0, 4.5 4.0 |
| | F | 2 | F | 1 | 7.5 | | | | | |
| <i>Prinia flaviventris</i> (R) | A | 2 | — | 4 | 7.0, 7.0, 7.2, 7.7 | | | | | |
| | K | 9 | — | 4 | 9.2, 9.8, 10.0, 10.4 | | | | | |
| <i>Prinia atrogularis</i> (R) | F | 1 | M | 1 | 6.9 | 8.9 ± 1.5 | — | I | + + + | 2.5 1.5 |
| | F | 2 | M | 1 | 7.2 | | | | | |
| | A | 1 | — | 4 | 9.2, 9.8, 10.4, 10.4 | | | | | |
| <i>Cisticola exilis</i> (R) | C | 4 | M | 1 | 11.5 | 10.2 ± 1.4 | — | I I 4 | — + + | 0.5 4.2 6.2/2.2, 5.0/1.2, 5.8/1.0, 5.9/— |
| | C | 1 | F | 1 | 8.0 | | | | | |
| | C | 4 | F | 3 | 10.0, 10.1, 11.2, — | | | | | |
| <i>Tesia olivea</i> (R) | F | 2 | M | 1 | 6.9 | — | — | I | — | (small) |
| | F | 2 | — | 1 | 6.3 | | | | | |
| <i>Cettia squameiceps</i> (M) | C | 1 | M | 1 | 7.9 | — | — | I | — | 1.0 |
| | C | 2 | M | 1 | 9.1 | | | | | |
| | E | 12 | — | 1 | 7.5 | | | | | |
| <i>Cettia fortipes</i> (M) | E | 12 | M | 1 | 7.3 | — | — | I | — | 3.8 |
| | F | 2 | M | 2 | 8.8, 10.5 | | | | | |
| <i>Bradypterus thoracicus</i> (M) | A | 2 | — | 1 | 9.3 | — | — | I | (small), (small) | |
| | H | 4 | M | 1 | 9.3 | | | | | |
| <i>Muscicapa sibirica</i> (M) | Ficedula parva | E | 12 | M | 1 | 9.0 | — | I | — | (small) |
| | D | 12 | — | 2 | 10.0, 10.8 | | | | | |
| <i>Ficedula strophiata</i> (M) | B | 2 | F | 1 | 12.4 | — | — | I | — | 4.7 |
| | B | 4 | F | 1 | 11.6 | | | | | |
| <i>Ficedula monileger</i> (R) | E | 12 | F | 1 | 11.0 | 11.0 ± 0.5 | — | 7 | — | 1.0, 1.5, 1.5, 2.0, 2.0, 2.0, 1.5 |
| | B | 4 | M | 4 | 10.7, 10.8, 11.1, 11.6 | | | | | |
| <i>Ficedula hyperythra</i> (R) | B | 1 | F | 4 | 10.5, 10.6, 10.6, 12.3 | 10.9 ± 0.8 | — | 4 | + | 6.0, 6.0, 6.2, 6.2 |
| | B | 4 | F | 1 | 10.7 | | | | | |
| <i>Ficedula hodgsonii</i> (M) | B | 1 | — | 3 | 10.5, 10.9, 13.2 | 10.5 ± 0.5 | — | I | + | 4.0/0.5 |
| | B | 4 | M | 2 | 8.3, 8.5 | | | | | |
| <i>Ficedula westermanni</i> (R) | B | 2 | M | 4 | 8.3, 8.4, 8.6, 8.8 | 8.5 ± 0.2 | — | 4 | — | 2.2, 1.4, 2.1, 1.1 |
| | E | 12 | F | 2 | 7.7, 8.2 | | | | | |
| <i>Cyanoptila cyanomelana</i> (M) | B | 1 | F | 2 | 7.8, 8.0 | 8.1 ± 0.3 | — | 2 | — | 3.5, 4.0 |
| | B | 2 | F | 1 | 8.3 | | | | | |
| <i>Niltava grandis</i> (R) | E | 5 | F | 1 | 8.5 | 8.5 ± 0.2 | — | I | — | 5.5 |
| | B | 1 | M | 1 | 10.4 | | | | | |
| <i>Niltava macgrigoriae</i> (R) | B | 4 | F | 1 | 10.2 | — | — | I | — | 3.0 |
| | E | 12 | M | 1 | 10.7 | | | | | |
| <i>Niltava sundara</i> (M) | B | 4 | M | 4 | 21.6, 21.8, 22.2, 22.5 | 22.7 ± 0.9 | — | 4 | — | 2.3, 2.5, 3.2, 4.3 (small), 1.5 |
| | E | 12 | M | 2 | 22.5, 22.7 | | | | | |
| <i>Niltava vivida</i> (M) | B | 1 | F | 3 | 22.5, 23.0, 24.2 | 22.7 ± 0.9 | — | 3 | — | 5.5, 4.5, 6.0 |
| | E | 12 | F | 1 | 21.7 | | | | | |
| <i>Cyornis banyumas</i> (R) | B | 1 | — | 1 | 24.5 | 24.5 ± 1.0 | — | I | — | 4.5 |
| | B | 2 | M | 1 | 31 | | | | | |
| <i>Cyornis rubeculoides glaucomans</i> (M) | B | 4 | — | 1 | 14.7 | 14.5 ± 1.0 | — | I | — | 1.5 |
| | B | 1 | M | 2 | 14.6, 15.3 | | | | | |
| <i>Cyornis banyumas</i> (R) | B | 1 | M | 1 | 12.8 | 14.5 ± 1.0 | — | I | — | 3.0, 3.0 |
| | E | 12 | M | 1 | 14.5 | | | | | |
| <i>Cyornis banyumas</i> (R) | B | 1 | F | 1 | 14.3 | 14.5 ± 1.0 | — | I | — | 4.5 |

| Species | Site | Month | Sex | n | Weights (g) | | mean | SD | n | Gonads | length (mm) |
|---|------|-------|-----|---|---|--|------|----|------|----------|--|
| | | | | | | | | | | activity | |
| <i>Culcicapa ceylonensis</i> (R) | B | 1 | M | 1 | 7.5 | | | | 1 | - | 1.5 |
| | B | 4 | M | 1 | 6.9 | | | | 1 | + | 5.3 |
| | B | 4 | F | 1 | 6.7 | | | | 1 | + | 4.4 |
| <i>Rhipidura hypoxantha</i> (R) | B | 2 | M | 1 | 5.2 | | | | 1 | + | 2.0 |
| | E | 12 | F | 1 | 4.4 | | | | 1 | - | 2.7 |
| <i>Rhipidura albicollis</i> | B | 1 | M | 2 | 10.0, 11.5 | | | | 2 | + | 3.0, 3.0 |
| | B | 1 | F | 5 | 9.8, 9.8, 10.1, 10.1, 10.2 | | | | 5 | + | 4.0/1.0, 4.5/-, 3.5/-, 5.5/1.0, 4.0/- |
| | B | 4 | F | 1 | 10.4 | | | | 1 | - | 4.0 |
| | E | 12 | F | 2 | 9.0, 9.2 | | | | | | (small), 4.5 |
| | B | 1 | - | 1 | 9.7 | | | | | | |
| <i>Rhipidura javanica</i> (R) | B | 4 | - | 2 | 11.2, 11.2 | | | | | | |
| | E | 12 | - | 4 | 10.0, 10.2, 10.5, 10.8 | | | | | | |
| | H | 4 | M | 1 | 12.5 | | | | 1 | + | 6.4 |
| | H | 4 | F | 2 | 12.5, 13.2 | | | | 2 | + | 4.5/1.5, 6.5/3.0 |
| <i>Hypothymis azurea</i> (R) | H | 4 | M | 2 | 13.1, 13.5 | | | | | | |
| | H | 9 | - | 8 | 10.7, 11.7, 11.9, 12.0, 12.0, 12.2, 13.8, 13.9 | | | | | | |
| | H | 9 | - | 1 | 10.9 | | | | | | |
| <i>Terpsiphone atrocaudata</i> (M) | H | 4 | M | 2 | 18.7 | | | | 1 | - | 2.5 |
| <i>Pachycephala cinerea</i> (R) | H | 4 | M | 2 | 17.1, 17.7 | | | | 2 | + | 8.0, 8.0 |
| <i>Motacilla cinerea</i> (M) | A | 2 | - | 1 | 16.2 | | | | | | |
| <i>Anthus hodgsoni</i> (M) | E | 12 | M | 2 | 21.0, 21.0 | | | | 2 | - | 1.5, 2.1 |
| | E | 12 | F | 1 | 21.6 | | | | 1 | - | 5.5 |
| | B | 4 | F | 2 | 18.9, 23.7 | | | | 2 | - | 5.0, 5.0 |
| <i>Anthus novaeseelandiae</i> <i>richardii</i> (M) | C | 1 | - | 1 | 29.6 | | | | | | |
| | (R) | A | 2 | - | 20.0 | | | | | | |
| <i>Anthus roseatus</i> (M) | A | 2 | - | 3 | 17.5, 22.4, 23.2 | | | | | | |
| <i>Lanius cristatus</i> (M) | D | 12 | - | 5 | 29.2, 30.3, 31.5, 31.5, 32.6 | | | | 31.5 | ± 1.7 | |
| | K | 9 | - | 1 | 34.1 | | | | | | |
| | H | 9 | - | 1 | 26.9 | | | | | | |
| <i>Lanius tigrinus</i> (M) | B | 4 | M | 1 | 40.5 | | | | 1 | + | 11.0 |
| | K | 9 | - | 1 | 46.1 | | | | | | |
| <i>Sturnus contra</i> (R) | H | 11 | - | 1 | 76.5 | | | | | | |
| <i>Aethopyga gouldiae</i> (M) | B | 1 | M | 1 | 6.9 | | | | 1 | - | 1.0 |
| | B | 1 | F | 1 | 6.1 | | | | 1 | - | 2.0 |
| <i>Aethopyga nipalensis</i> (R) | E | 4 | M | 1 | 6.3 | | | | 1 | + | 4.1 |
| | E | 12 | M | 3 | 6.0, 6.3, 6.5 | | | | 3 | - | (all small) |
| | E | 12 | F | 1 | 5.4 | | | | 1 | - | (n.m.) |
| | E | 5 | - | 1 | 5.6 | | | | | | |
| <i>Aethopyga saturata</i> (R) | B | 4 | M | 2 | 5.5, 5.7 | | | | 2 | + | 5.0, 5.0 |
| | E | 12 | M | 1 | 5.7 | | | | 1 | - | (small) |
| | B | 1 | F | 2 | 4.6, 5.2 | | | | 2 | + | 3.1/-, 5.5/1.0 |
| | B | 4 | F | 1 | 5.0 | | | | 1 | + | 2.9/1.0 |
| <i>Arachnothera magna</i> (R) | E | 12 | M | 1 | 32.9 | | | | 1 | - | 2.0 |
| <i>Zosterops erythopleura</i> (M) | B | 1 | M | 1 | 11.5 | | | | 1 | - | 1.0 |
| | C | 4 | M | 2 | 9.9, 11.2 | | | | 2 | - | 1.0, 1.4 |
| <i>Zosterops japonica</i> (M) | C | 1 | M | | 10.2 | | | | 1 | - | 1.0 |
| | E | 12 | M | 2 | 9.4, 9.5 | | | | 2 | - | 1.0, 1.3 |
| | E | 12 | F | 6 | 9.0, 9.1, 9.2, 9.3, 9.8, 9.8 | | | | 6 | - | 4.5, (small), 5.0, 5.0, 5.0, (small) |
| | E | 12 | - | 4 | 8.8, 9.3, 9.5, 9.8 | | | | | | |
| <i>Zosterops palpebrosa</i> (R) | B | 4 | M | 1 | 7.2 | | | | 1 | + | 3.7 |
| <i>Passer flaveolus</i> (R) | H | 4 | - | 1 | 16.6 | | | | | | |
| <i>Ploceus manyar</i> (R) | L | 5 | M | 1 | 18.4 | | | | 1 | + | 3.5 |
| | L | 5 | M | 1 | 16.4 | | | | 1 | - | 1.5 |
| <i>Ploceus hypoxanthus</i> (R) | F | 1 | M | 1 | 18.6 | | | | 1 | - | (small) |
| <i>Lonchura striata</i> (R) | F | 2 | - | 1 | 11.5 | | | | | | |
| <i>Lonchura punctulata</i> (R) | F | 2 | - | 4 | 11.6, 12.0, 12.3, 12.3 | | | | | | |
| | A | 2 | - | 1 | 13.8 | | | | | | |
| | K | 9 | - | 2 | 12.8, 12.9 | | | | | | |
| | D | 12 | - | 1 | 13.0 | | | | | | |
| <i>Emberiza fucata</i> (M) | A | 2 | - | 2 | 21.1, 22.5 | | | | | | |
| <i>Emberiza aureola</i> (M) | A | 2 | - | 1 | 17.8 | | | | | | |
| <i>Emberiza rutila</i> (M) | B | 2 | M | 2 | 15.2, 16.2 | | | | 2 | - | 1.3, 1.2 |
| | B | 1 | F | 1 | 15.8 | | | | 1 | - | 3.5 |

Acknowledgements. We thank R. J. Dobias, R. Eve, Ms. V. J. Hamilton, Ms. A-M Guigue, G. C. Yong, Tawee Gorsatah, Werasak Intajom and Vandee Nongngork for their assistance in collection of data. Dr. Rauf Ali provided statistical advice. The Royal Thai Forest Department kindly provided assistance and accommodation at Doi Pha Hom Pok, Doi Inthanon, Bang Phra and Khao Sam Roi Yot and the Fisheries Division of the Department of Agriculture at Beung Boraphet and Samut Sakhon. Most of the data were collected on expeditions funded by the Smithsonian Institution. We are also very grateful to Dr. Boonsong Lekagul for both logistic and financial support.

References:

- von Bröckel, K. 1973. Vergleichende Messungen an lebenden und frischtoten Gartengrasmücken (*Sylvia borin*). *J. Orn.* 114: 114-122.
- Chasen, F. N. & Kloss, C. B. 1932. On birds from Doi Sutep, 5,600 feet, north Siam. *J. Siam Soc. Nat. Hist. Suppl.* 8: 231-248.
- Deignan, H. G. 1945. The birds of northern Thailand. *U.S. Natn. Mus. Bull.* 186: 1-616.
- Herbert, E. G. 1923-26. Nests and eggs of birds in Central Siam. *J. Nat. Hist. Soc. Siam* 6: 81-123, 215-222, 293-322, 323-356.
- McClure, H. E. 1974a. *Migration and Survival of the Birds of Asia*. U.S. Army Medical Component, SEATO Medical Research Laboratory, Bangkok.
- McClure, H. E. 1974b. Some bionomics of the birds of Khao Yai National Park, Thailand. *Nat. Hist. Bull. Siam Soc.* 25: 99-194.
- McClure, H. E. & Kwanyuen, P. 1973. The avifaunal complex of an Open-billed Stork Colony (*Anastomus oscitans*) in Thailand. *Nat. Hist. Bull. Siam Soc.* 25: 133-156.
- Medway, Lord. 1973. A ringing study of migratory Barn Swallows in West Malaysia. *Ibis* 115: 60-85.
- Nisbet, I. C. T. 1967. Migration and moult in Pallas's Grasshopper Warbler. *Bird Study* 14: 96-103.
- Nisbet, I. C. T. & Medway, Lord. 1972. Dispersion, population ecology and migration of Eastern Great Reed Warblers *Acrocephalus orientalis* wintering in Malaysia. *Ibis* 114: 451-494.
- Riley, J. H. 1938. Birds from Siam and the Malay Peninsula in the United States National Museum collected by Drs. Hugh M. Smith and William L. Abbott. *U.S. Natn. Mus. Bull.* 172: 1-581.
- Round, P. D. 1982. Notes on breeding birds in north-west Thailand. *Nat. Hist. Bull. Siam. Soc.* 30: 1-14.
- Sokal, R. R. & Rohlf, F. J. 1981. *Biometry*, 2nd edition. W. H. Freeman & Co., San Francisco.
- Ward, P. 1969. The annual cycle of the Yellow-vented Bulbul *Pycnonotus goiavier* in a humid equatorial environment. *J. Zool. Soc. Lond.* 156: 25-46.
- Addresses:* D. S. Melville and P. D. Round, Association for the Conservation of Wildlife, 4 Old Custom House Lane, Bangkok 10500, Thailand. (Present address for D. S. Melville: WWF Hong Kong, GPO Box 12721, Hong Kong).

© British Ornithologists' Club 1984.

The Rufous Sparrows of the Cape Verde Islands

by D. Summers-Smith

Received 22 February 1984

The African Rufous Sparrows occur in a number of widely separated populations south of 18°N. They were first described by A. Smith (1836) as *Pyrgita Motitensis*, with the type locality subsequently indentified by Winterbottom (1966) as Motito in northern Cape Province, South Africa. (*Pyrgita* is now recognised as a synonym for *Passer* and has been suppressed.) Darwin (1841) collected a male from S. Tiago island in the Cape Verde Archipelago in 1832 during the voyage of the *Beagle*; this was later given the name *Pyrgita iagoensis* in 1837 by Gould (1837).